

# The love for the three conics

Stefan Liebscher and Dierck-E. Liebscher

## Abstract

Neville's theorem of the three focus-sharing conics finds a simplification and a new outreach in the context of projective geometry.

## 1 Introduction

In Euclidean geometry, let three points in the plane serve as three pairs of foci for three conics. The three pairs of conics define three lines through their intersections, and the three lines are concurrent, they pass a common point [1, 2, 3].

This theorem has found devotees and different proofs in Euclidean geometry [4, 5, 6]. It may be expanded to non-Euclidean geometries (Fig. 1) where it allows a generalising formulation in projective terms [7].

From the perspective of projective geometry, foci are defined for pairs of conics [8]. They are the vertices of the quadrilateral of common tangents and come in three pairs, which define





Figure 1: Three focus-sharing conics in Cayley-Klein geometries.



The pencil can be seen as determined by the four common points, or as generated by two conics with their four intersections. Selecting any of the conics as absolute conic (strong-line ellipse), the pencil is determined by any other conic, in particular the singular conic given by a pair of focal lines.

Figure 2: Pencil of conics through four common points.


the three diagonals of the quadrilateral [9]. Linear combination of the two conics yields the pencil of conics tangent to all four sides of the quadrilateral.

For a single conic, it is the metric-generating absolute conic of Cayley-Klein geometries which may serve as the second conic to define foci in the usual sense. This is the case in the Neville problem. Confocal conics conventionally form the pencil defined by two foci and the absolute conic in the background. Pencils exist for each pair of the three focus-sharing conics, too. It is the consideration of these pencils which admits the short and snappy proofs (see ch. 9 in [7]).

## 2 Pairs of conics

To explore the relation between two conics, we consider the generated pencils of conics. Figure 2 shows the pencil with the four intersection points fixed. It contains three singular elements: the three pairs of common chords. Figure 3 shows the dual pencil with the four common tangents fixed. It also contains three singular elements: the three pairs of foci. To emphasize their duality, we call the common chords also focal lines. Whichever focus we choose, it determines two directrices, which intersect in a diagonal point common to both the quadrilateral of tangents and the quadrangle of common points (Fig. 4). This diagonal point carries one of the three pairs of focal lines (i.e. common chords), see Prop. 1 below.

For Neville's problem, we start with three points (Fig. 5). For any two of them we choose



The dual pencil can be seen as determined by the four common tangents, or as generated by two conics with their four common tangents. Selecting any of the conics as absolute conic (strong-line ellipse), the pencil is determined by any other conic, in particular the singular conic given by a pair of foci. The plane contains points where two conics of the dual pencil intersect, and points which meet none of the conics.


Figure 3: Dual pencil of conics with four common tangents.

a conic for which the two points are a pair of foci with respect to the absolute conic. Any two of the conics share just one focus with one crossing of directrices and one singular conic consisting of a pair of common chords. We obtain such a pair of common chords for each of the three pairs of conics. The condition that the shared foci are also foci with respect to the absolute conic imposes linear dependence, and the common chords are the edges of a quadrangle of points in which the common chords intersect. This turns out to be simple algebra, and yields the desired proofs.

We use homogeneous coordinates and write “ $\sim$ ” to denote equality up to a nonzero scalar factor. Conics can be represented by pairs  $[\mathbf{c}, \mathbf{C}]$  of symmetric matrices. They provide a linear map of points onto lines:  $Q \mapsto \mathbf{c}Q$ , the polar of  $Q$ , and its dual, a linear map of lines onto points,  $g \mapsto \mathbf{C}g$ , the pole of  $g$ . The pair of both maps is also called polarity. The peripheral points of a conic are the zeros of the quadratic form,  $P^T \mathbf{c} P = 0$ , its tangents are the zeros of the dual,  $t^T \mathbf{C} t = 0$ .

For regular conics, the two matrices  $\mathbf{C}$  and  $\mathbf{c}$  are reciprocal,  $\mathbf{C}\mathbf{c} \sim \mathbf{1}$ . Singular conics satisfy  $\mathbf{C}\mathbf{c} = 0 = \mathbf{c}\mathbf{C}$ . If one of  $\mathbf{c}, \mathbf{C}$  has at least rank 2, the other one is (a scalar multiple of) its adjoint (or transpose cofactor) matrix.

For  $\mathbf{c}$  of rank 2, we can write  $\mathbf{c} = gh^T + hg^T$ . This yields a singular conic consisting of two distinct lines  $g, h$  of peripheral points with a pencil of tangents through the double point  $g \times h$ . For  $\mathbf{C}$  of rank 2, we can write  $\mathbf{C} = PQ^T + QP^T$  and find a singular conic consisting of a (double) line  $P \times Q$  of peripheral points through the concurrency centers  $P, Q$  of the two



Two conics have 4 common tangents and 4 common points. On the left, part of the pencil with the common points is shown, with the six chords (focal lines) and the three diagonal points. On the right, part of the pencil with the common tangents is shown, with the six intersections (foci) and the three diagonal lines. The diagonal triangles coincide.

Figure 4: Two pencils for a pair of conics with their coinciding diagonal triangles.


pencils of tangents.

For two distinct regular conics  $[k_1, K_1], [k_2, K_2]$ , the pencil  $\mathcal{P}[k_1, k_2]$  is given by the linear combinations  $\{\alpha_1 k_1 + \alpha_2 k_2\}$  (Fig. 2), the dual pencil  $\tilde{\mathcal{P}}[K_1, K_2]$  by  $\{\beta_1 K_1 + \beta_2 K_2\}$  (Fig. 3).

The pencil is uniquely determined by its four common points,  $\mathcal{P} = \mathcal{P}[Q_1, \dots, Q_4]$ . Its singular elements are the pairs of opposite chords, i.e. the pairs of opposite edges of the quadrangle  $Q_1, \dots, Q_4$  of common points intersecting in the diagonal points of the quadrangle. We note that for regular real conics the singular elements are real even in the case of only complex intersections of the conics.

Analogously, the dual pencil is uniquely determined by the quadrilateral of common tangents,  $\tilde{\mathcal{P}} = \tilde{\mathcal{P}}[t_1, \dots, t_4]$ . Its singular elements are the diagonal lines of the quadrilateral with tangent pencils in the vertices of the quadrilateral. The six intersections of the tangents  $t_i$ , the vertices of the quadrilateral, are the common foci of the dual pencil  $\tilde{\mathcal{P}}$ .

**Proposition 1** *For any pair of conics, the triangle given by the diagonal points of the quadrangle of common points coincides with the triangle given by the diagonal lines of the quadrilateral*



In the case of three ellipses, three of the common chords have no real intersection points, nevertheless the chords are real, and the six chords are edges of a quadrangle of intersections. In the case of three hyperbolas, we have chosen the eccentricities large enough to obtain only real intersections of the conics and real chords. For clearness, both drawings show an euclidean setting, for the general setting, see Fig. 1.

Figure 5: Three focus-sharing conics with the quadrangles of Neville centers.

of common tangents. This diagonal triangle is also self-polar: Each diagonal point is the pole of the opposite diagonal line with respect to all conics of the two pencils.

**Proof.** See Fig. 4 and Thm. 7.6 of [7].  $\diamond$

In other words: given a pair of conics, the diagonals of the tangent quadrilateral meet in the three intersection points of opposite common chords.

### 3 Three focus-sharing conics

In Neville's problem, three conics  $\mathbf{K}_k$ ,  $k = 1, 2, 3$  share three foci  $F_k$ ,  $k = 1, 2, 3$  with respect to the absolute conic  $\mathbf{C}$ . The pair  $[F_l, F_m]$  belongs to  $\mathbf{K}_k$  ( $k, l, m$  cyclically), and  $F_k$  is focus in the pairs  $[\mathbf{K}_l, \mathbf{K}_m]$ ,  $[\mathbf{K}_l, \mathbf{C}]$ , and  $[\mathbf{K}_m, \mathbf{C}]$ .

The pair  $[\mathbf{K}_l, \mathbf{K}_m]$  itself has six foci (again in three pairs).  $F_k$  is one of them, and its adjoint partner on the diagonal passing  $F_k$  will be denoted by  $F_k^*$ . The line  $d_k \sim F_k \times F_k^*$  is a diagonal

line of the quadrilateral of tangents common to  $\mathbf{K}_l$  and  $\mathbf{K}_m$ . The opposite diagonal point  $D_k$  can be represented as pole of  $d_k$  to both of the conics,  $D_k \sim \mathbf{K}_l d_k \sim \mathbf{K}_m d_k$ , and the four polars  $\mathbf{k}_l F_k, \mathbf{k}_m F_k, \mathbf{k}_l F_k^*, \mathbf{k}_m F_k^*$  are incident with  $D_k$ .

$$D_k \sim \mathbf{K}_l d_k \sim \mathbf{K}_m d_k \sim \mathbf{k}_l F_k^* \times \mathbf{k}_m F_k. \quad (1)$$

The focus  $F_k$  has a directrix (its polar) with both  $\mathbf{K}_l$  and  $\mathbf{K}_m$ , and the intersection of the two is the diagonal point  $D_k$  (Fig. 5).

We now use the identity of the diagonal triangles, Prop. 1. The diagonal point  $D_k$  is incident with two opposite chords of the pencil  $\mathcal{P}(\mathbf{k}_l, \mathbf{k}_m)$ . They form one of the singular elements of this pencil,  $\mathbf{n}_k$ . We can write

$$\mathbf{n}_k \sim (D_k^T \mathbf{k}_l D_k) \mathbf{k}_m - (D_k^T \mathbf{k}_m D_k) \mathbf{k}_l. \quad (2)$$

After substituting one  $D_k$  by  $\mathbf{K}_l d_k$ , the other by  $\mathbf{K}_m d_k$ , we obtain

$$\mathbf{n}_k \sim (d_k^T \mathbf{K}_m \mathbf{k}_l \mathbf{K}_l d_k) \mathbf{k}_m - (d_k^T \mathbf{K}_l \mathbf{k}_m \mathbf{K}_m d_k) \mathbf{k}_l \sim (d_k^T \mathbf{K}_m d_k) \mathbf{k}_m - (d_k^T \mathbf{K}_l d_k) \mathbf{k}_l \quad (3)$$

This is a singular conic consisting of two lines, i.e. a pair of common chords, and a double point, the diagonal point  $D_k$ . For each pair  $[\mathbf{K}_l, \mathbf{K}_m]$  of the three conics, we obtain such a singular conic consisting of that pair of opposite common chords which intersect in the diagonal point  $D_k$ .

**Proposition 2** *The three singular conics  $\mathbf{n}_k$  are linear dependent. Therefore, the common chords represented in these singular conics are the six sides of a quadrangle (Fig. 5).*

**Proof.** The conics  $\mathbf{K}_m$  are elements of the dual pencils generated by  $\mathbf{C}$  and the singular conics  $F_k F_l^T + F_l F_k^T$ . We fix representatives to remove the ambiguity of arbitrary nonzero scalar factors and write


$$\mathbf{K}_m = \omega_m \mathbf{C} + \lambda_m (F_k F_l^T + F_l F_k^T). \quad (4)$$

Since the diagonal  $d_m$  is incident with  $F_m$ , we obtain

$$(d_k^T \mathbf{K}_m d_k) = \omega_m (d_k^T \mathbf{C} d_k). \quad (5)$$

Equation (3) now yields  $\mathbf{n}_{lm} = \omega_m \mathbf{k}_m - \omega_l \mathbf{k}_l$  (again choosing a suitable representative to remove the nonzero coefficient  $d_k^T \mathbf{C} d_k$ ) and linear dependency  $\mathbf{n}_{12} + \mathbf{n}_{23} + \mathbf{n}_{31} = 0$  emerges.  $\bowtie$

Starting with the absolute conic  $\mathbf{C}$ , we can reformulate Prop. 2.



The figure shows three conics  $\mathbf{K}_m$  outside the absolute conic  $\mathbf{C}$  in order to get real lines and points. The points  $F_m$  are the shared foci. The points  $F_m^*$  are opposite to the foci  $F_m$  in the quadrilateral of tangents common to  $\mathbf{K}_k$  and  $\mathbf{K}_l$ . They are incident on a line.

Figure 6: Three focus-sharing conics with their collinear adjoint foci.

**Corollary 3** *Given a conic with three pairs of tangents defining three intersections. Given one representative of each dual pencil generated by two of the three pairs of tangents. Then, the three intersections have two polars each (w.r.t. the conics of the associated dual pencils). These polars intersect in points which carry two common chords. The six common chords are edges of a quadrangle.*


We turn again to the adjoint foci. A pair  $[\mathbf{K}_l, \mathbf{K}_m]$  defines six foci. They yield three diagonals which define a pairing of the foci, and the  $[F_k, F_k^*]$  is such a pair. The focus  $F_k$  is the intersection of two tangents common to  $[\mathbf{K}_l, \mathbf{K}_m]$ , and  $F_k^*$  is the intersection of the two other common tangents. The focus  $F_k$  is a focus of the pair  $[\mathbf{C}, \mathbf{K}_l]$  and of the pair  $[\mathbf{C}, \mathbf{K}_m]$  as well. The focus adjoint to  $F_k$  in the pair  $[\mathbf{K}_l, \mathbf{K}_m]$  is  $F_k^*$ , in the pair  $[\mathbf{C}, \mathbf{K}_l]$  it is  $F_m$ , and in the pair  $[\mathbf{C}, \mathbf{K}_m]$  it is  $F_l$ .

The following proposition was proven by Bogdanov [4] in the Euclidean case. In Cayley-Klein geometries, it is a generalization of the four-conics theorem known as the dual of the four-conics-theorem (see Fig. 17 in [10]).

**Proposition 4** *The three adjoint foci of the three focus-sharing conics are collinear. The six foci  $F_k, F_k^*, i = 1 \dots 3$ , are the vertices of a quadrilateral (Fig. 6).*

**Proof.** We refer again the singular members of dual pencils. The singular elements of the dual pencil  $\mathbf{K}_{lm} = \alpha \mathbf{K}_l + \beta \mathbf{K}_m$  are given by the diagonal points itself:

$$\mathbf{N}_k = (d_k^T \mathbf{K}_l d_k) \mathbf{K}_m - (d_k^T \mathbf{K}_m d_k) \mathbf{K}_l. \quad (6)$$



Given three shared foci and three ellipses, one adds the six directrices first. We choose one of the two triangles contained in the convex hexagon ( $d_{13}, d_{21}, d_{32}$ ) to establish trilinear coordinates  $\xi_k$ . The distance of foci ( $2f_i$ ) is found in the distance of the directrices ( $2\kappa_i = 2f_i/\varepsilon_i^2$ ), so that intersections of two ellipses are given by comparing the distances from the focus (e.g.  $F_1$ ) with the distances from the directrices ( $\xi_2$  and  $2\kappa_3 - \xi_3$  for the common chord of  $\mathbf{K}_2$  and  $\mathbf{K}_3$ ).

Figure 7: Three focus-sharing ellipses on the Euclidean plane.

We take (5) and obtain

$$\sum_{k \in \{1,2,3\}} \omega_k (d_l^T \mathbf{C} d_l) (d_m^T \mathbf{C} d_m) \mathbf{N}_k = 0. \quad (7)$$

The three singular conics (all are pairs  $F_k F_k^*$  of foci) are linear dependent, i.e. the foci  $F_k, F_k^*$ ,  $k = 1, 2, 3$  are the vertices of a complete quadrilateral, the three adjoint foci  $F_k^*$  are collinear.  $\bowtie$

## 4 The limit of the Neville points

Any point can be found as a Neville point, if it is reached by the three pencils of conics generated by the three pairs of foci with the absolute conic. In particular, any point in the inner part of the triangle can be the intersection of three ellipses of the set.

There is one particular point among the Neville centers. It is the limit for collapsing ellipses, for the singular members of the three dual pencils  $\tilde{\mathcal{P}}[\mathbf{K}_m, \mathbf{C}]$ . This limit is given by  $\omega_m \rightarrow 0$  in eq. (4). We start with the Euclidean picture of Neville's proof (Fig 7). The triangle of foci is augmented with the six directrices. We identify two triangles of three directrices each. They are congruent and similar to the triangle of foci. We use distances  $(\xi_1, \xi_2, \xi_3)$  to the sides of one of them ( $d_{13}, d_{21}, d_{32}$ ). After the calculation,  $[\xi_1 : \xi_2 : \xi_3]$  can be reinterpreted as trilinear coordinates. We denote the eccentricity of the conic  $\mathbf{K}_k$  by  $\varepsilon_k$ , the distances of the foci by  $2f_k$ , and the distance of the directrices by  $2\kappa_k = 2f_k/\varepsilon_k^2$ . The common chords of the three ellipses

are

$$\varepsilon_1 \xi_1 = \varepsilon_2 (2\kappa_2 - \xi_2), \quad \varepsilon_2 \xi_2 = \varepsilon_3 (2\kappa_3 - \xi_3), \quad \varepsilon_3 \xi_3 = \varepsilon_1 (2\kappa_1 - \xi_1), \quad (8)$$

We obtain the intersection

$$\xi_1^S = \kappa_1 + \varepsilon_1^{-1}(\kappa_2 \varepsilon_2 - \kappa_3 \varepsilon_3), \quad \xi_2^S = \kappa_2 + \varepsilon_2^{-1}(\kappa_3 \varepsilon_3 - \kappa_1 \varepsilon_1), \quad \xi_3^S = \kappa_3 + \varepsilon_3^{-1}(\kappa_1 \varepsilon_1 - \kappa_2 \varepsilon_2). \quad (9)$$

The limit of collapsing ellipses is simply given by eccentricities equal one. In this case, the equations (8) for the chords reduce to the equations for the angular bisectors of the directrices. The common chords become the angular bisectors of the triangle  $[D_1, D_2, D_3]$  approaching the focus triangle, too. The intersection of the chords approaches the incenter for eccentricities equal one.

We shall show, using the singular elements of pencils of conics, that in the general Cayley-Klein geometry, the limit of the intersection again approaches the incenter. More specifically, we shall show that the limit of the common chord of two conics sharing a focus is the angular bisector. We show the proof for the case of a regular absolute conic.

The shared foci are  $F_k, F_l$  of  $[\mathbf{K}_m, \mathbf{C}]$ , the diagonal points  $D_m \sim \mathbf{k}_k F_m \times \mathbf{k}_l F_m \sim \mathbf{K}_k d_m \sim \mathbf{K}_l d_m$ . The focal lines (i.e. the common chords) are the lines of the singular elements  $\mathbf{n}_m \sim (D_m^\top \mathbf{k}_l D_m) \mathbf{k}_k - (D_m^\top \mathbf{k}_k D_m) \mathbf{k}_l$ . As we are interested in the limit  $\omega_m \rightarrow 0$ , we fix  $\lambda_m = 1$  in (4) and express the dual conics as

$$\mathbf{K}_m = \omega_m \mathbf{C} + F_k F_l^\top + F_l F_k^\top. \quad (10)$$

**Proposition 5** *Given the dual conics (10). Then the the singular conics  $\mathbf{n}_m$ , given by (3), of the pencils  $\mathcal{P}(\mathbf{k}_k, \mathbf{k}_l)$  are formed by the angular bisectors*

$$\begin{aligned} \mathbf{w}_m \sim & ((F_m \times F_k)^\top \mathbf{C} (F_m \times F_k)) (F_m \times F_l) (F_m \times F_l)^\top \\ & - ((F_m \times F_l)^\top \mathbf{C} (F_m \times F_l)) (F_m \times F_k) (F_m \times F_k)^\top, \end{aligned} \quad (11)$$

i.e.  $\mathbf{n}_m \sim \mathbf{w}_m$ .

**Proof.** *Step 1:  $\mathbf{k}_m$  as inverse of  $\mathbf{K}_m$  with suitable scaling.* We normalize  $F_k$  such that  $\gamma_k := F_k^\top \mathbf{c} F_k = \pm 1$  and write  $p_k := \mathbf{c} F_k$ ,  $\delta_m := F_k^\top \mathbf{c} F_l = F_k^\top p_l = F_l^\top p_k$ . The coefficient  $\omega_m$  in (10) is determined now. The limit of eccentricity 1 means  $\omega_m \rightarrow 0$  here. The adjoint  $\mathbf{k}_m$  to  $\mathbf{K}_m$  is found as

$$\mathbf{k}_m = \omega_m \mathbf{K}_m^{-1} = \mathbf{c} + \frac{1}{(\mu_m^2 - \gamma_k \gamma_l)} (\gamma_l p_k p_k^\top + \gamma_k p_l p_l^\top) - \frac{\mu_m}{(\mu_m^2 - \gamma_k \gamma_l)} (p_l p_k^\top + p_k p_l^\top) \quad (12)$$

with  $\mu_m = \omega_m + \delta_m$ .

*Step 2: Singular conics  $\mathbf{n}_m$  of common chords.* We use (3) and (5) from the proof of Prop. 2 to find

$$\mathbf{n}_m \sim \mathbf{k}_k - \mathbf{k}_l. \quad (13)$$

*Step 3:  $\mathbf{k}_m$  in the limit of eccentricity 1.* We fix

$$\tilde{\mathbf{k}}_m = (F_k \times F_l)(F_k \times F_l)^\top \quad (14)$$

as representation of the singular conic given by the line through the foci  $F_k, F_l$ . Equation (12) yields

$$F_k^\top \mathbf{k}_m F_k = \frac{\gamma_k(\mu_m - \delta_m)^2}{\mu_m^2 - \gamma_k \gamma_l}, \quad F_l^\top \mathbf{k}_m F_l \text{ the same,}$$

and finally

$$F_k^\top \mathbf{k}_m F_l = \frac{(\mu_m - \delta_m)(\mu_m \delta_m - \gamma_k \gamma_l)}{\mu_m^2 - \gamma_k \gamma_l}.$$

In the limit  $\omega_m \rightarrow 0$ , that is  $\mu_m \rightarrow \delta_m$ , these values vanish. Then both  $F_k$  and  $F_l$  are incident with the  $(\mathbf{k}_m)$ -polars of  $F_k$  and  $F_l$ . Consequently, the polar of any point on  $F_k \times F_l$  is this line itself:

$$\lim_{\omega_m \rightarrow 0} \mathbf{k}_m \sim \tilde{\mathbf{k}}_m \quad (15)$$

*Step 4: Comparing coefficients.* At this point, we only have to check that the coefficients of  $\mathbf{k}_k$  and  $\mathbf{k}_l$  in (13) fit the coefficients in (11):

$$\lim_{\omega \rightarrow 0} (\mathbf{k}_k - \mathbf{k}_l) \stackrel{?}{\sim} ((F_m \times F_k)^\top \mathbf{C}(F_m \times F_k)) \tilde{\mathbf{k}}_k - ((F_m \times F_l)^\top \mathbf{C}(F_m \times F_l)) \tilde{\mathbf{k}}_l.$$


Because of (15), it suffices to check one regular point. We choose  $Q = \mathbf{C}(F_k \times F_l)$  and obtain

$$Q^\top \tilde{\mathbf{k}}_m Q = ((F_k \times F_l)^\top Q)^2, \quad Q^\top \mathbf{k}_m Q = (F_k \times F_l)^\top Q$$

Therefore, indeed,

$$\begin{aligned} \lim_{\omega \rightarrow 0} \mathbf{k}_k - \mathbf{k}_l &= \frac{1}{(F_m \times F_l)^\top \mathbf{C}(F_m \times F_l)} \tilde{\mathbf{k}}_k - \frac{1}{(F_m \times F_k)^\top \mathbf{C}(F_m \times F_k)} \tilde{\mathbf{k}}_l \\ &\sim ((F_m \times F_k)^\top \mathbf{C}(F_m \times F_k)) \tilde{\mathbf{k}}_k - ((F_m \times F_l)^\top \mathbf{C}(F_m \times F_l)) \tilde{\mathbf{k}}_l. \end{aligned}$$

The singular conic of the diagonals is the singular conic of the angular bisectors, and the Neville center coincides with the incenter of the focus triangle.  $\bowtie$



Each pair of foci generates a quadrilateral of tangents to the absolute conic. There are two other pairs of intersections, which are connected by the other two diagonals of the quadrilateral. These six diagonals are again edges of a quadrangle: It is the quadrangle of circumcenters of the focus triangle.

Figure 8: The quadrangle of diagonals of the set of the three pairs of foci.

## 5 Trifles

In the Euclidean setup, eq. (8), we obtain an incenter for the Neville point also in the case of equal eccentricities. The incenter is not that of the focus triangle, but of the directrix hexagon.

In the limit of infinite eccentricity, the Neville point approaches the circumcenter of the focus triangle. This circumcenter is one of the 60 Brianchon points of the tangent hexagon, which is generated by the three foci. More general: To each pair of foci belong two other pairs corresponding to the two other diagonals. These six diagonals are again edges of a quadrangle (Fig. 8).

The trilinear coordinates of the incenter of the triangle  $[D_1, D_2, D_3]$  relative to the triangle  $[U_1, U_2, U_3]$  in Fig. 7 are given by


$$N = [(a + b - c) : (b + c - a) : (c + a - b)]. \quad (16)$$

This seems to be a particular point [11] in the triangle  $[U_1, U_2, U_3]$ , but it comes in 32 different versions: The hexagon of directrices admits eight choices of directrix triangles, and for each triangle a quadrangle of in- and excenters:

$$N = [(\pm a + s_\gamma b - s_\beta c) : (\pm b + s_\alpha c - s_\gamma a) : (\pm c + s_\beta a - s_\alpha b)] \quad (17)$$

with  $s_\alpha, s_\beta, s_\gamma = \pm 1$  and  $s_\alpha s_\beta s_\gamma = 1$  for the quadrangle.

Since we calculate in projective spaces, the propositions of section 3 have dual counterparts. Instead of foci as intersections of tangents common with a fourth (absolute) conic, we



Dual to the focus-sharing conics we obtain intersections of common tangents in a quadrilateral and connections of focal lines and their adjoints as edges of a quadrangle.

Figure 9: Three conics sharing focal lines.

consider the chords through intersections with this fourth (absolute) conic. We also call them focal lines to emphasize this duality. Three pairs of tangents to the absolute conic generated three dual pencils from which three conics were taken. Now three pairs of points on the absolute conic generate three pencils from which three conics can be taken. Each pair of the three conics had four intersections and three pairs of chords. Now each pair of three conics has four common tangents and three pairs of foci. In each pair of conics, one pair of chords intersected in the intersection of the polars of the shared focus. Now in each pair of conics, one pair of foci is collinear with the poles of the shared focal lines. The three chosen pairs of chords were the edges of a quadrangle. Now the three chosen pairs of foci are the vertices of a quadrilateral. This is the dual version of Prop. 2 (Fig. 9, left).

Each pair of three conics sharing a focal line defines an adjoint line (the partner in the pair to which the shared focal line belongs). Together with the shared focal lines we obtain six lines which are found to be the edges of a complete quadrangle. This is the dual version of Prop. 4 (Fig. 9, right), known in Euclidean notions as the four-conics theorem [10].

We conclude with a general remark on the relation between Euclidean and non-Euclidean constructions. Theorems about conics and lines only, without explicit reference to symmetry and perpendiculars, can be interpreted as theorems in non-Euclidean geometry. Figure 6 is an




Figure 10: Collinearity between quadrilaterals of common tangents.

example. One may state that if two common tangents of each pair of three conics touch a fourth conic, then the remaining common tangents of each pair intersect in three collinear points [10]. This formulation is purely Euclidean, but misses the non-Euclidean connection. In addition, Euclidean theorems about lines and circles should be expected to find an simple non-Euclidean extension by use of the non-Euclidean definition of circles and perpendicularity. Figure 10 shows the connection between Monge's theorem, a dual three-conic theorem cited in [10, 12], and a Pascal-line construction cited in [6].

## References

- [1] Eric Harold Neville. A focus-sharing set of three conics. *Math. Gaz.*, 20(239):182–183, 1936.
- [2] Eric Harold Neville. The focus-sharing conics again. *Math. Gaz.*, 21(242):56, 1937.
- [3] Lawrence Crawford. Determination of the foci, directrices, axes and eccentricities of a conic whose equation is given with numerical coefficients. *Math. Gaz.*, 18:43–46, 1934.
- [4] Ilya I. Bogdanov. Two theorems on the focus-sharing ellipses: a three-dimensional view. *J. Class. Geom.*, 1:1–5, 2012.

- [5] A.V. Akopyan and A.A. Zaslavsky. *Geometry of conics.*, volume 26 of *Mathematical world*. Providence, RI: American Mathematical Society (AMS), 2007.
- [6] Alexander Skutin. Some new theorems in plane geometry. *arXiv:1704.04923v3 math.MG*, pages 1–19, 2017.
- [7] Stefan Liebscher. *Projektive Geometrie der Ebene. Ein klassischer Zugang mit interaktiver Visualisierung*. Heidelberg: Springer Spektrum, 2017.
- [8] Jürgen Richter-Gebert. *Perspectives on projective geometry. A guided tour through real and complex geometry*. Springer, Berlin, 2011.
- [9] Stefan Liebscher and Dierck-E. Liebscher. The relativity of conics and circles. *Forum Geom.*, 18:1–6, 2018.
- [10] Evelyn C.J.A. Money-Coutts G.B. and Tyrrell J.A. *The seven circles theorem and other new theorems*. London : Stacey, 1974.
- [11] Clark Kimberling. Encyclopedia of triangle centers. <http://faculty.evansville.edu/ck6/>.
- [12] Arseniy Akopyan. On some classical constructions extended to hyperbolic geometry. *arXiv:1105.2153v1 math.MG*, pages 1–12, 2011.

Stefan Liebscher  
 TNG Technology Consulting GmbH  
 Unterföhring  
 contact@stefan-liebscher.de  
<https://www.stefan-liebscher.de/geometry>

Dierck-E. Liebscher  
 Leibniz-Institut für Astrophysik  
 Potsdam  
 deliebscher@aip.de  
<http://www.dierck-e-liebscher.de>